

# A NON SMOOTH EXPONENTIAL\*

Esteban Andruchow

Instituto de Ciencias – Univ. Nac. de Gral. Sarmiento  
Argentina

## Abstract

Let  $\mathcal{M}$  be a type  $\text{II}_1$  von Neumann algebra,  $\tau$  a trace in  $\mathcal{M}$ , and  $L^2(\mathcal{M}, \tau)$  the GNS Hilbert space of  $\tau$ . If  $L^2(\mathcal{M}, \tau)_+$  is the completion of the set  $\mathcal{M}_{sa}$  of selfadjoint elements, then each element  $\xi \in L^2(\mathcal{M}, \tau)_+$  gives rise to a selfadjoint unbounded operator  $L_\xi$  on  $L^2(\mathcal{M}, \tau)$ . In this note we show that the exponential  $\exp : L^2(\mathcal{M}, \tau)_+ \rightarrow L^2(\mathcal{M}, \tau)$ ,  $\exp(\xi) = e^{iL_\xi}$  is continuous but non differentiable. The same holds for the Cayley transform  $C(\xi) = (L_\xi - i)(L_\xi + i)^{-1}$ . We also show that the unitary group  $U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  with the strong operator topology, is not an embedded submanifold of  $L^2(\mathcal{M}, \tau)$ , in any way which makes the product  $(u, w) \mapsto uw$  ( $u, w \in U_{\mathcal{M}}$ ) a differentiable map.

**Keywords:** unitary group, non commutative integration.

## 1 Introduction

Let  $\mathcal{M}$  be a type  $\text{II}_1$  von Neumann algebra with a faithful and normal tracial state  $\tau$ . Let  $L^2(\mathcal{M}, \tau)$  be the Hilbert space obtained by completion of  $\mathcal{M}$  with the norm  $\|x\|_2 = \tau(x^*x)^{1/2}$ . By Segal's theory of abstract integration [3], any element  $\xi \in L^2(\mathcal{M}, \tau)$  can be regarded as a (possibly unbounded) operator  $L_\xi$  on  $L^2(\mathcal{M}, \tau)$ , affiliated to  $\mathcal{M}$ , as follows (see [1]). Let  $J$  be the antiunitary involution of  $L^2(\mathcal{M}, \tau)$ , which on the dense subspace  $\mathcal{M} \subset L^2(\mathcal{M}, \tau)$  is just the involution  $*$  of  $\mathcal{M}$ , and for  $m \in \mathcal{M}$  consider the linear map  $m \mapsto Jm^*J\xi$ . This map is a closable operator, and  $L_\xi$  is its closure.

The elements  $m \in \mathcal{M}$  will be considered as operators acting by left multiplication on  $L^2(\mathcal{M}, \tau)$ , when regarded as *elements* of  $L^2(\mathcal{M}, \tau)$ , they will be denoted by  $\vec{m}$ .

An interesting fact [3] is that if  $\xi$  satisfies  $J\xi = \xi$ , then the associated operator  $L_\xi$  is selfadjoint. Let us denote by  $L^2(\mathcal{M}, \tau)_+ = \{\xi \in L^2(\mathcal{M}, \tau) : J\xi = \xi\}$ . Clearly  $L^2(\mathcal{M}, \tau)_+$  is a real Hilbert space, and the inner product of  $L^2(\mathcal{M}, \tau)$  is real and symmetric when restricted to  $L^2(\mathcal{M}, \tau)_+$ . Indeed,  $L^2(\mathcal{M}, \tau)_+$  is the completion of the set  $\mathcal{M}_{sa}$  of selfadjoint elements of  $\mathcal{M}$ , and if  $\vec{m}_1, \vec{m}_2 \in \mathcal{M}_{sa}$ , then  $\langle \vec{m}_1, \vec{m}_2 \rangle = \tau(m_2 m_1) = \tau(m_1 m_2) = \langle \vec{m}_2, \vec{m}_1 \rangle$ .

In this note we consider the map

$$\exp : L^2(\mathcal{M}, \tau)_+ \rightarrow U_{\mathcal{M}} \vec{1} \subset L^2(\mathcal{M}, \tau), \quad \exp(\xi) = e^{iL_\xi} \vec{1},$$

---

\*2000 Mathematics Subject Classification: 46L51, 58B10, 58B25.

where  $U_{\mathcal{M}}$  is the unitary group of  $\mathcal{M}$ , and  $U_{\mathcal{M}\vec{1}}$  is just the same set regarded as a subset of  $L^2(\mathcal{M}, \tau)$ , which induces on  $U_{\mathcal{M}}$  a metric topology equivalent to the strong operator topology. In what follows we identify  $U_{\mathcal{M}}$  and  $U_{\mathcal{M}\vec{1}}$ .

We prove that the map  $\exp$  is continuous but non smooth, in fact, non differentiable. We consider the Cayley transform

$$C : L^2(\mathcal{M}, \tau)_+ \rightarrow U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau), \quad C(\xi) = (L_{\xi} - i)(L_{\xi} + i)^{-1}\vec{1},$$

which is also continuous and non differentiable.

The unitary group  $U_{\mathcal{M}}$  in the strong operator topology can be embedded in  $L^2(\mathcal{M}, \tau)$  as a complete topological group. The group operations are clearly continuous in the  $L^2$ -metric, and  $U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  is closed. We finish this note by proving that it cannot be embedded as a differentiable Banach-Lie group.

## 2 Non regularity of $\exp$ and $C$

Let us see first that these maps are continuous. The following lemma will be useful. It relates the  $L^2$  topology with the generalization of the strong topology to unbounded operators. Our reference on this subject is [2].

**Lemma 2.1** *If a sequence  $\xi_n$  converges in  $L^2(\mathcal{M}, \tau)_+$  to  $\xi$ , then the operators  $L_{\xi_n}$  converge to  $L_{\xi}$  in the strong resolvent sense.*

**Proof.** Suppose that  $\xi_n \in L^2(\mathcal{M}, \tau)_+$  converges to  $\xi$ . Then for  $m \in \mathcal{M}$ ,  $L_{\xi_n}m\vec{1}$  converges to  $L_{\xi}m\vec{1}$ . Indeed,  $L_{\xi_n}m\vec{1} = Jm^*J\xi_n \rightarrow Jm^*J\xi$  because  $Jm^*J$  is bounded, and this last vector equals  $L_{\xi}m\vec{1}$ . We claim that  $\mathcal{M}\vec{1}$  is a common core for all (selfadjoint) operators of the form  $L_{\eta}$ ,  $\eta \in L^2(\mathcal{M}, \tau)_+$ . In that case, using VIII.25 of [2] it follows that  $L_{\xi_n}$  converges to  $L_{\xi}$  in the strong resolvent sense. Our claim follows using another result in [2], VIII.11: If  $A$  is a selfadjoint operator and  $D \subset D(A)$  is a dense subspace which is invariant by the one parameter group  $e^{itA}$ , i.e.  $e^{itA}(D) \subset D$  for all  $t \in \mathbb{R}$ , then  $D$  is a core for  $A$ . Now clearly  $e^{itL_{\eta}}m\vec{1} \in \mathcal{M}\vec{1}$ , because  $e^{itL_{\eta}} \in \mathcal{M}$  for all  $t$  if  $\eta \in L^2(\mathcal{M}, \tau)_+$ . It follows that  $\mathcal{M}\vec{1}$  is a core for  $L_{\eta}$ .  $\square$

It shall be useful to have an alternative writing for  $C$ . Note that  $(L_{\xi} - i)^{-1}$  and  $(L_{\xi} + i)^{-1}$  commute, and that  $(L_{\xi} - i)^{-1}(\xi - i\vec{1}) = \vec{1}$ . Therefore  $(L_{\xi} + i)^{-1}\vec{1} = (L_{\xi} - i)^{-1}(L_{\xi} + i)^{-1}(\xi - i\vec{1})$ , and then

$$C(\xi) = (L_{\xi} + i)^{-1}(\xi - i\vec{1}), \quad \xi \in L^2(\mathcal{M}, \tau)_+.$$

**Proposition 2.2** *The maps  $\exp$  and  $C$  are continuous.*

**Proof.** If  $\xi_n \rightarrow \xi$  in  $L^2(\mathcal{M}, \tau)_+$ , then the resolvents  $(L_{\xi_n} + i)^{-1}$  converge strongly to the resolvent  $(L_{\xi} + i)^{-1}$ . Note also that these operators are contractions. On the other hand  $\xi_n - i\vec{1} \rightarrow \xi - i\vec{1}$  in  $L^2(\mathcal{M}, \tau)_+$ . It follows that  $C(\xi_n) = (L_{\xi_n} - i)^{-1}(\xi_n - i\vec{1})$  converge to  $C(\xi)$ . The same type of argument shows that the function  $\exp$  is continuous. Indeed, if  $L_{\xi_n}$  converges to  $L_{\xi}$  in the strong resolvent sense, and  $f$  is a bounded continuous function on  $\mathbb{R}$ , then  $f(L_{\xi_n}) \rightarrow f(L_{\xi})$  strongly ([2], VII.20). Therefore  $\exp(\xi) = e^{iL_{\xi}}\vec{1}$  is continuous.  $\square$

Although these maps are not differentiable, they do have directional derivatives at the origin.

**Lemma 2.3** *For all  $\eta, v \in L^2(\mathcal{M}, \tau)_+$ , the curve  $t \mapsto C(\eta + tv)$  is differentiable at  $t = 0$ , and*

$$\frac{d}{dt}C(\eta + tv)|_{t=0} = \frac{dC}{dv}(\eta) = -2i(L_{\eta} + i)^{-1}J(L_{\eta} - i)^{-1}Jv.$$

**Proof.** Note that

$$C(\eta + tv) - C(\eta) = (L_{\eta+tv} + i)^{-1}(\eta + tv - i\vec{1}) - (L_{\eta} + i)^{-1}(\eta - i\vec{1}).$$

The above sum can be decomposed in the following terms:

$$(L_{\eta+tv} + i)^{-1}(\eta + tv - i\vec{1}) - (L_{\eta+tv} + i)^{-1}(\eta - i\vec{1})$$

and

$$(L_{\eta+tv} + i)^{-1}(\eta - i\vec{1}) - (L_{\eta} + i)^{-1}(\eta - i\vec{1}).$$

We deal first with the first term, which equals

$$(L_{\eta+tv} + i)^{-1}(tv) = t(L_{\eta+tv} + i)^{-1}(v).$$

The second term equals

$$((L_{\eta+tv} + i)^{-1} - (L_{\eta} + i)^{-1})(\eta - i\vec{1}) = (L_{\eta+tv} + i)^{-1}[L_{\eta} + i - (L_{\eta+tv} + i)](L_{\eta} + i)^{-1}(\eta - i\vec{1}).$$

Note that this expression is well defined, since the vector  $(L_{\eta} + i)^{-1}(\eta - i\vec{1}) = C(\eta) \in \mathcal{M}\vec{1}$  lies in the domain of any combination of the operators  $L_{\nu}$ ,  $\nu \in L^2(\mathcal{M}, \tau)_+$ . Moreover, it apparently equals

$$-t(L_{\eta+tv} + i)^{-1}L_{\nu}(L_{\eta} + i)^{-1}(L_{\eta} - i)\vec{1} = t(L_{\eta+tv} + i)^{-1}J(L_{\eta} + i)(L_{\eta} - i)^{-1}Jv.$$

Putting both terms together one has

$$\frac{C(\eta + tv) - C(\eta)}{t} = (L_{\eta+tv} + i)^{-1}(v - J(L_{\eta} + i)(L_{\eta} - i)^{-1}Jv).$$

If we let  $t$  tend to 0, then  $\eta + tv \rightarrow \eta$  in  $L^2(\mathcal{M}, \tau)_+$  and  $(L_{\eta+tv} + i)^{-1} \rightarrow (L_{\eta} + i)^{-1}$  strongly. Therefore the derivative of  $C(\eta + tv)$  at  $t = 0$  exists and equals

$$(L_{\eta+v} + i)^{-1}(v - J(L_{\eta} + i)(L_{\eta} - i)^{-1}Jv).$$

Finally, the vector  $v - J(L_{\eta} + i)(L_{\eta} - i)^{-1}Jv$  can be written

$$v - J(L_{\eta} + i)(L_{\eta} - i)^{-1}Jv = J(1 - (L_{\eta} + i)(L_{\eta} - i)^{-1})Jv = -2iJ(L_{\eta} - i)^{-1}Jv.$$

If one replaces this last expression in the result obtained for the derivative, one obtains the desired formula.  $\square$

**Lemma 2.4** *For any  $v \in L^2(\mathcal{M}, \tau)_+$ , the curve  $t \mapsto \exp(tv)$  is differentiable at  $t = 0$  and*

$$\frac{d}{dt}\exp(tv)|_{t=0} = \frac{d}{dv}\exp(0) = iv.$$

**Proof.** The one parameter unitary group  $t \mapsto e^{itL_v}$  is strongly differentiable at  $t = 0$  on the domain of  $L_v$  [2], i.e. if  $\xi \in D(L_v)$ ,  $\frac{d}{dt}e^{itL_v}\xi|_{t=0}$  exists and equals  $iL_v\xi$ . Taking  $\xi = \vec{1} \in D(L_v)$  proves the result.  $\square$

**Theorem 2.5** *The maps  $\exp : L^2(\mathcal{M}, \tau)_+ \rightarrow U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  and  $C : L^2(\mathcal{M}, \tau)_+ \rightarrow U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  are not differentiable on any neighbourhood of  $0 \in L^2(\mathcal{M}, \tau)_+$ .*

**Proof** Suppose that  $exp$  is differentiable on a neighbourhood  $0 \in \mathcal{V} \subset L^2(\mathcal{M}, \tau)_+$ . For any  $\xi \in L^2(\mathcal{M}, \tau)$  let  $\xi = \xi_+ + \xi_-$  be the decomposition of  $\xi$  in  $L^2(\mathcal{M}, \tau) = L^2(\mathcal{M}, \tau)_+ \oplus L^2(\mathcal{M}, \tau)_-$ . We shall construct a local diffeomorphism on  $L^2(\mathcal{M}, \tau)$ , which restricted to  $L^2(\mathcal{M}, \tau)_+$  will provide a local homeomorphism onto a neighbourhood of  $\vec{1}$  in  $U_{\mathcal{M}}$ . Afterwards we shall prove that this fact leads to contradiction. Consider the map

$$\theta : L^2(\mathcal{M}, \tau) \rightarrow L^2(\mathcal{M}, \tau), \quad \theta(\xi) = exp(\xi_+) + i\xi_-.$$

The projections  $\xi \mapsto \xi_+$  and  $\xi \mapsto \xi_-$  are (real) linear and bounded, therefore they are  $C^\infty$ . It follows that  $\theta$  is differentiable in  $\mathcal{V}$ . Note that  $\theta(0) = \vec{1}$  and  $\frac{d}{d\xi}exp(0) = i\xi$ . Therefore

$$d\theta_0(\xi) = i\xi_+ + i\xi_- = i\xi.$$

By the inverse function theorem it follows that there exists a ball  $0 \in \mathcal{B}_\epsilon(0) \subset \mathcal{V}$  in the  $\|\cdot\|_2$ -metric and an open set  $\mathcal{W}$  of  $L^2(\mathcal{M}, \tau)$  such that  $\theta : \mathcal{B}_\epsilon(0) \rightarrow \mathcal{W}$  is a diffeomorphism. Note that  $\theta$  maps  $\mathcal{B}_\epsilon(0) \cap L^2(\mathcal{M}, \tau)_+$  onto  $\mathcal{W} \cap U_{\mathcal{M}}$ , i.e.  $\theta|_{L^2(\mathcal{M}, \tau)_+}$  is a local homeomorphism between  $\mathcal{B}_\epsilon(0)$  and a neighbourhood of  $\vec{1}$  in  $U_{\mathcal{M}}$  in the  $L^2$ -topology.

Fix  $\delta^{1/2} < \epsilon$ , and for each integer  $n \geq 1$  choose a projection  $p_n \in \mathcal{M}$  such that  $\tau(p_n) = \delta/n^2$ . Put  $a_n = np_n$ . Note that  $\|a_n\|_2 = \delta^{1/2}$ . Indeed,  $\tau(a_n^*a_n) = n^2\tau(p_n) = \delta$ . Therefore  $a_n \in \mathcal{B}_\epsilon(0)$  and  $a_n$  does not tend to 0. On the other hand

$$\|exp(a_n) - 1\|_2^2 = 2 - \tau(e^{ia_n}) - \tau(e^{-ia_n}).$$

Clearly

$$\tau(e^{ia_n}) = 1 + \sum_{k \geq 1} \tau\left(\frac{(in)^k}{k!} p_n\right) = 1 + \frac{\delta}{n^2}(e^{in} - 1).$$

Analogously  $\tau(e^{-ia_n}) = 1 + \frac{\delta}{n^2}(e^{-in} - 1)$ . It follows that  $exp(a_n) \rightarrow \vec{1}$  but  $\theta^{-1}(exp(a_n))$  does not tend to 0, a contradiction.

To prove the same result for  $C$ , one proceeds analogously. Using the fact that if  $C$  were differentiable, then by 2.3,  $dC_0(\xi) = -2i\xi$ , one can construct also in this case a local homeomorphism between a ball centered at  $0 \in L^2(\mathcal{M}, \tau)_+$  and a  $L^2$ -neighbourhood of  $-\vec{1}$  in  $U_{\mathcal{M}}$  (note that  $C(0) = -\vec{1}$ ). Let  $p_n \neq 0$  be projections in  $\mathcal{M}$  such that  $\tau(p_n) \rightarrow 0$ . Then, as above,  $\|1 - exp(p_n)\|_2 \rightarrow 0$ . Note also that  $1 - exp(p_n)$  has non trivial kernel, indeed,  $1 - exp(p_n) = (e^i - 1)p_n$ . On the other hand, if  $C$  were a local homeomorphism, then there would be a neighbourhood  $-1 \in \mathcal{U} \subset U_{\mathcal{M}}$  where all  $v \in \mathcal{U}$  would satisfy that  $v - 1$  has trivial kernel, because unitaries in the range of the Cayley transform have this property.  $\square$

For the Cayley transform one has the following weaker regularity conditions.

**Proposition 2.6** *The Cayley transform  $C$  is weakly  $C^1$ , i.e. for any fixed  $\nu \in L^2(\mathcal{M}, \tau)$ , the complex valued map  $\xi \mapsto \langle C(\xi), \nu \rangle$  is  $C^1$ .*

*If we regard  $C$  as a map from  $L^2(\mathcal{M}, \tau)_+$  to  $L^1(\mathcal{M}, \tau)$ , it is differentiable.*

**Proof.** For any  $\eta, v \in L^2(\mathcal{M}, \tau)_+$ , using 2.3 one has

$$\frac{d}{dv} \langle C, \nu \rangle (\eta) = \langle \frac{dC}{dv}(\eta), \nu \rangle = -2i \langle (L_\eta + i)^{-1} J(L_\eta - i)^{-1} Jv, \nu \rangle.$$

Recall that if  $\eta_n \rightarrow \eta$  in  $L^2(\mathcal{M}, \tau)_+$ , then the resolvents  $(L_{\eta_n} - i)^{-1}$  and  $(L_{\eta_n} + i)^{-1}$  are contractions which converge strongly to  $(L_\eta - i)^{-1}$  and  $(L_\eta + i)^{-1}$ . It follows that  $\frac{d}{dv} \langle C, \nu \rangle (\eta)$  is continuous in both parameters  $\nu, v \in L^2(\mathcal{M}, \tau)_+$ , and  $C$  is weakly  $C^1$ . Let us prove that

$$C : L^2(\mathcal{M}, \tau)_+ \rightarrow L^1(\mathcal{M}, \tau)$$

is differentiable. Using the computations done in 2.3, one has that

$$C(\eta + v) - C(\eta) = -2i(L_{\eta+v} + i)^{-1}J(L_{\eta} - i)^{-1}Jv$$

and

$$\frac{dC}{dv}(\eta) = -2i(L_{\eta} + i)^{-1}J(L_{\eta} - i)^{-1}Jv$$

therefore  $C(\eta + v) - C(\eta) - \frac{dC}{dv}(\eta)v$  equals

$$-2i[(L_{\eta+v} + i)^{-1} - (L_{\eta} + i)^{-1}]J(L_{\eta} - i)^{-1}Jv.$$

We must show that the norm 1 of this expression divided by  $\|v\|_2$  tends to zero if  $v$  tends to zero in  $L^2(\mathcal{M}, \tau)$ . Denote by  $\Delta = (L_{\eta+v} + i)^{-1} - (L_{\eta} + i)^{-1}$  and by  $\psi = J(L_{\eta} - i)^{-1}Jv$ . Note that  $\Delta \in \mathcal{M}$  and  $\psi \in L^2(\mathcal{M}, \tau)$  with  $\|\psi\|_2 \leq \|v\|_2$ . Also

$$\|\Delta\psi\|_1 \leq \|\Delta\|_2\|\psi\|_2.$$

Indeed, let  $\vec{x}_n$  be a sequence in  $\mathcal{M}\vec{1}$  converging to  $\psi$  in  $L^2(\mathcal{M}, \tau)$ . Then  $\|\Delta\vec{x}_n\|_1 = \tau(|\Delta x_n|) = \tau(u^*\Delta x_n)$  where  $u$  is the partial isometry in the polar decomposition of  $\Delta x_n \in \mathcal{M}$ , which can be chosen unitary since  $\mathcal{M}$  is finite. By the Cauchy Schwarz inequality  $\tau(u\Delta x_n) \leq \tau(\Delta^*\Delta)^{1/2}\tau(x_n^*x_n)^{1/2}$ . Since  $x_n \rightarrow \psi$ ,  $\Delta x_n \rightarrow \Delta\psi$  in  $L^2(\mathcal{M}, \tau)$ , and the inequality follows. Therefore

$$\frac{\|C(\eta + v) - C(\eta) - \frac{dC}{dv}(\eta)v\|_1}{\|v\|_2} = 2\frac{\|\Delta\psi\|_1}{\|v\|_2} \leq 2\|\Delta\|_2.$$

The proof ends by showing that  $\|\Delta\|_2 \rightarrow 0$  as  $v$  tends to zero. Clearly  $\Delta$  tends to zero in the strong operator topology, because  $L_{\eta+v} \rightarrow L_{\eta}$  in the strong resolvent sense. Then

$$\tau(\Delta^*\Delta) = \langle \Delta\vec{1}, \Delta\vec{1} \rangle \rightarrow 0.$$

□

We now state the result of the non embedddability of  $U_{\mathcal{M}}$  in  $L^2(\mathcal{M}, \tau)$  as a Lie group.

**Theorem 2.7** *The unitary group  $U_{\mathcal{M}}$  of  $M$ , with the  $L^2$  metric is not an embedded submanifold of  $L^2(\mathcal{M}, \tau)$  with differentiable multiplication map  $(u, w) \mapsto uw$  ( $u, w \in U_{\mathcal{M}}$ ).*

**Proof.** The proof consists of showing that if  $U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  were an embedded submanifold, then it would be a Lie group, with Lie algebra identified with  $L^2(\mathcal{M}, \tau)_- := \{\xi \in L^2(\mathcal{M}, \tau) : J\xi = -\xi\}$ . Moreover, the Lie bracket would extend the commutator of (antselfadjoint) elements of  $\mathcal{M}$ ,  $[x, y] = xy - yx$ . This is clearly not possible, the commutators of elements of  $L^2(\mathcal{M}, \tau)$  lie in  $L^1(\mathcal{M}, \tau)$ , eventually outside of  $L^2(\mathcal{M}, \tau)$  [3].

Suppose that  $U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  is a submanifold. If  $u(t)$  is a smooth curve of unitaries with  $u(0) = 1$  and  $u'(0) = \xi$ , differentiating  $u^*(t)u(t) = 1$  at  $t = 0$  yields  $J\xi + \xi = 0$ , i.e.  $\xi \in L^2(\mathcal{M}, \tau)_-$ . Also any element  $\xi \in L^2(\mathcal{M}, \tau)_-$  can be obtained as the velocity vector of a curve in  $U_{\mathcal{M}}$ . It was shown above that the curve  $u(t) = \exp(tv)$  is differentiable at  $t = 0$  for any  $v \in L^2(\mathcal{M}, \tau)_+$ , put  $v = -i\xi$ , then  $u'(0) = \xi$ . The tangent space of  $U_{\mathcal{M}}$  at a point  $u \in U_{\mathcal{M}}$  clearly identifies with  $uL^2(\mathcal{M}, \tau)_-$ .

The multiplication is differentiable by hypothesis. The inversion  $u \mapsto u^{-1} = u^*$  is continuous ( $\mathcal{M}$  is finite). It can be regarded as the restriction of a real linear map of  $L^2(\mathcal{M}, \tau)$ , namely  $J$ , and therefore is differentiable.

Therefore  $U_{\mathcal{M}}$  is a Lie group, and its Lie algebra identifies with  $L^2(\mathcal{M}, \tau)_-$ . Let us compute the bracket under this identification. The left action of the group  $U_{\mathcal{M}}$  on itself,  $\ell^u : U_{\mathcal{M}} \rightarrow U_{\mathcal{M}}$ ,

$\ell^u(w) = uw$  extends to a linear bounded operator on  $L^2(\mathcal{M}, \tau)$ . Then if  $\xi \in L^2(\mathcal{M}, \tau)_-$ , the left invariant vector field induced by  $\xi$  is  $X_\xi(u) = u\xi$  ( $u \in U_{\mathcal{M}}$ ). If  $f$  is a smooth function on a neighbourhood of  $\vec{1} \in U_{\mathcal{M}}$ , then the derivative  $X_\xi f$  can be computed as follows:

$$X_\xi f(u) = \frac{d}{dt} f(ue^{tL\xi})|_{t=0} = df_u(u\xi),$$

where  $df_u$  is the tangent map of  $f$  at  $u \in U_{\mathcal{M}}$ . Note that in the above computation again one only uses that  $t \mapsto e^{tL\xi}\vec{1}$  is differentiable at  $t = 0$  ( $\xi \in L^2(\mathcal{M}, \tau)_-$ ). Let  $\vec{x}, \vec{y} \in i\mathcal{M}_{sa} \subset L^2(\mathcal{M}, \tau)_-$  two antiselfadjoint elements on  $\mathcal{M}$  (note that  $i\mathcal{M}_{sa}$  is dense in  $L^2(\mathcal{M}, \tau)_-$ ). Let us compute  $X_{\vec{x}}X_{\vec{y}}f$ :

$$X_{\vec{x}}X_{\vec{y}}f(u) = \frac{d}{dt} df_{ue^{tx}}(ue^{tx}\vec{y})|_{t=0} = d^2f_u(u\vec{x}, u\vec{y}) + df_u(u\vec{x}\vec{y}).$$

Since  $d^2f_u$  is a symmetric bilinear form, it follows that the bracket  $[X_{\vec{x}}, X_{\vec{y}}]$  is given by

$$[X_{\vec{x}}, X_{\vec{y}}]f(u) = df_u(u\vec{x}\vec{y} - u\vec{y}\vec{x}) = df_u(u(xy - \vec{y}x)),$$

which coincides with the left invariant derivation  $X_{xy - \vec{y}x}f(u)$ . This says that the bracket of  $x, y$  regarded as an element of the Lie algebra of  $U_{\mathcal{M}}$  is the usual commutator  $xy - yx$ .  $\square$

It would be interesting to know if the result holds dropping the assumption on the differentiability of the multiplication. That is, if  $U_{\mathcal{M}} \subset L^2(\mathcal{M}, \tau)$  is never an embedded submanifold.

## References

- [1] E. Christensen, Universally bounded operators on von Neumann algebras of type II<sub>1</sub>, Algebraic methods in operator theory, 195-204, Birkäuser Boston, Boston, MA, 1994.
- [2] M. Read, B. Simon, Methods of Modern Mathematical Physics, vol. I: Functional Analysis, Academic Press, New York, 1978.
- [3] I.E. Segal, A non commutative extension of abstract integration, Ann. Math. 57 (1953), 401-457.

Esteban Andruchow  
 Instituto de Ciencias  
 Universidad Nacional de Gral. Sarmiento  
 J. M. Gutierrez entre J.L. Suarez y Verdi  
 (1613) Los Polvorines  
 Argentina  
 e-mail: eandruch@ungs.edu.ar